arXiv:hep-th/9302039v1 10 Feb 1993

NBI-HE-93-08
February 1993

Free Field Realization of $N = 2$ Super W_3 Algebra

Katsushi Ito*

Niels Bohr Institute, Blegdamsvej 17, DK-2100, Copenhagen Ø, Denmark

Abstract

We study the quantum $N = 2$ super-W_3 algebra using the free field realization, which is obtained from the supersymmetric Miura transformation associated with the Lie superalgebra $A(2|1)$. We compute the full operator product expansions of the algebra explicitly. It is found that the results agree with those obtained by the OPE method.

---

*Address after March 1, 1993: Institute of Physics, University of Tsukuba, Ibaraki 305, Japan
E-mail: ito@nbivax.nbi.dk

W-algebraic extensions of the $N = 2$ superconformal algebra [1] have attracted much interest in the context of conformal field theories[2, 3], topological W-gravities[4] and more recently in non-critical W-strings[5]. There are at least two approaches to study quantum W-algebras[6]. One is based on the OPE method[7], in which W-algebras can be constructed by requiring the associativity and closure of the algebra. The other approach is the free field realization[8], in which the generators are expressed by free fields through the quantum Miura transformations. In this approach the closure of the algebra is not obvious at quantum level, in particular for the W-algebras associated with non simply laced Lie algebra[9].

In the case of $N = 2$ super-W algebras, it has been understood that the free field realization can be obtained from the quantum hamiltonian reduction [10] of affine Lie superalgebras $A(n|n-1)^{(1)}$ [2, 3]. The classical Poisson brackets structure in the case of $A(2|1)^{(1)}$ has been studied by using techniques such as the super Gel'fand-Dickii bracket [11] and Polyakov's soldering procedure [12, 13]. In the quantum case, the full algebra has been presented by using the OPE method [14, 15]. In the free field approach, however, only some of the operator product expansions have been calculated [3].

In this letter, we study the free field realization of $N = 2$ super-W_3 algebra and compute the full operator product expansions. We find that the present free field approach gives the same algebra as the one obtained by the OPE method.

We begin with the supersymmetric Miura transformation based on a Lie superalgebra $A(n|n-1) = sl(n+1|n)$. We take the purely odd simple root system $\{\alpha_1, \ldots, \alpha_{2n}\}$, with the Cartan matrix $\alpha_i \cdot \alpha_j = (-1)^{i-1}\delta_{i+1,j}$. Let $\{\lambda_1, \ldots, \lambda_{2n}\}$ be the fundamental weights of $A(n|n-1)$ satisfying $\alpha_i \cdot \lambda_j = \delta_{ij}$. Denote $N = 1$ super-holomorphic coordinate by $Z = (z, \theta)$ and a super-derivative by $D = \frac{\partial}{\partial\theta} + \theta\partial$ ($\partial \equiv \frac{\partial}{\partial z}$). We introduce $2n$ free bosons $\phi(z) = (\phi_1(z), \ldots, \phi_{2n}(z))$ and real fermions $\psi(z) = (\psi_1(z), \ldots, \psi_{2n}(z))$ satisfying the operator product expansions (OPEs):

$$\phi_i(z)\phi_j(w) = -\delta_{ij}\ln(z-w) + \dots, \qquad \psi_i(z)\psi_j(w) = \frac{\delta_{ij}}{z-w} + \dots\ . \tag{1}$$

Define the scalar superfields by $\Phi(Z) = \phi(z) + \mathrm{i}\theta\psi(z)$. The $A(n|n-1)$ super Miura transformation [16] is defined as

$$D^{2n+1} + \sum_{i=2}^{2n+1} W_{\frac{i}{2}}(Z)(aD)^{2n+1-i} =: (aD-\Theta_{2n+1}(Z))(aD-\Theta_{2n}(Z))\cdots(aD-\Theta_1(Z)) :, \tag{2}$$

where $\Theta_i(Z) = (-1)^{i-1}(\lambda_i - \lambda_{i-1})\cdot D\Phi(Z)$ $(\lambda_0 = \lambda_{2n+1} \equiv 0)$ and $a = -\mathrm{i}\alpha_+$. The symbol $:\ :$ denotes the normal ordering. The currents $W_{i/2}(Z)$ $(i = 2, \dots, 2n+1)$ generate the $N = 2$ super W_{n+1} algebra. In the following we consider the case $n = 2$ and use the component formalism. Let us define bosonic and fermionic W-currents as follows:

$$\begin{aligned} W_1(Z) &= J(z) + \mathrm{i}\theta[G^+(z) + G^-(z)], \\ W_{\frac{3}{2}}(Z) &= a[\mathrm{i}G^-(z) + \theta T(z)], \\ W_2(Z) &= W_2(z) + \mathrm{i}\theta[W^+(z) + W^-(z)], \\ W_{\frac{5}{2}}(Z) &= a[\mathrm{i}W^-(z) + \theta W_3(z)]. \end{aligned} \tag{3}$$

It is convenient to introduce complex bosons $\varphi_i(z), \bar{\varphi}_i(z)$ $(i = 1, 2)$ and complex fermions $\chi_i(z)$, $\bar{\chi}_i(z)$ $(i = 1, 2)$ by $\varphi_i = \lambda_{2i}\cdot\phi(z)$, $\bar{\varphi}_i = \alpha_{2i}\cdot\phi(z)$, $\chi_i = \lambda_{2i}\cdot\psi(z)$ and $\bar{\chi}_i = \alpha_{2i}\cdot\psi(z)$. From the Miura transformation (2), we find that the generators of the $N = 2$ super-W_3 algebra are given by

$$\begin{aligned} J &= U_1 + U_2 - a\partial\bar{\varphi}_2, \\ G^+ &= G_1^+ + G_2^+, \\ G^- &= G_1^- + G_2^- - a\partial\bar{\chi}_2, \\ T &= T_1 + T_2 - a\partial^2\bar{\varphi}_2, \\ W_2 &= a^2\partial U_1 - a\bar{\chi}_2 G_1^+ + U_1 U_2 - a^3\partial^2\bar{\varphi}_2 - a\partial\bar{\varphi}_2 U_2, \\ W^+ &= a^2\partial G_1^+ + a\partial\bar{\varphi}_2 G_1^+ + U_2 G_1^+ + U_1 G_2^+ - a\partial\bar{\varphi}_2 G_2^+, \end{aligned}$$

$$\begin{aligned}
W^- &= a^2\partial G_1^- - a^3\partial^2\bar{\chi}_2 + U_2 G_1^- + U_1 G_2^- + a\bar{\chi}_2 T_1 - a\bar{\chi}_2\partial U_1 - aU_2\partial\bar{\chi}_2 - a\partial\bar{\varphi}_2 G_2^-, \\
W_3 &= a^2\partial T_1 + U_2 T_1 + T_2 U_1 - G_2^+ G_1^- - G_1^+ G_2^- + a\partial\bar{\varphi}_2 T_1 - a\partial\bar{\varphi}_2 T_2 \\
&\quad -a\partial\bar{\varphi}_2\partial U_1 - a\bar{\chi}_2\partial G_1^+ - a\partial\bar{\chi}_2 G_2^+ - a\partial^2\bar{\varphi}_2 U_2 - a^3\partial^3\bar{\varphi}_2,
\end{aligned} \tag{4}$$

where $U_i = \chi_i\bar{\chi}_i - a\partial(\bar{\varphi}_i - \varphi_i)$, $G_i^+ = \chi_i\partial\bar{\varphi}_i + a\partial\chi_i$, $G_i^- = -\bar{\chi}_i\partial\varphi_i - a\partial\bar{\chi}_i$ and $T_i = -\partial\varphi_i\partial\bar{\varphi}_i - \bar{\chi}_i\partial\chi_i - a\partial^2\bar{\varphi}_i$ for $i = 1, 2$. $T(z)$ denotes the twisted energy-momentum tensor: $T(z) = T^{N=2}(z) + \frac{1}{2}\partial J(z)$, where $T^{N=2}(z)$ is the energy-momentum tensor of the $N = 2$ model with central charge $c^{N=2} = 6(1 + 3a^2)$. The OPEs of the twisted $N = 2$ algebra are given by:

$$\begin{aligned}
T(z)T(w) &= \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w} + \ldots, \\
T(z)J(w) &= \frac{-2(1+3a^2)}{(z-w)^3} + \frac{J(w)}{(z-w)^2} + \frac{\partial J(w)}{z-w} + \ldots, \\
T(z)G^+(w) &= \frac{G^+(w)}{(z-w)^2} + \frac{\partial G^+(w)}{z-w} + \ldots, \quad T(z)G^-(w) = \frac{2G^-(w)}{(z-w)^2} + \frac{\partial G^-(w)}{z-w} + \ldots, \\
J(z)J(w) &= \frac{2(1+3a^2)}{(z-w)^2} + \ldots, \quad J(z)G^{\pm}(w) = \frac{\pm G^{\pm}(w)}{z-w} + \ldots, \\
G^+(z)G^-(w) &= \frac{2(1+3a^2)}{(z-w)^3} + \frac{J(w)}{(z-w)^2} + \frac{T(w)}{z-w} + \ldots, \quad G^{\pm}(z)G^{\pm}(w) = \text{regular}.
\end{aligned} \tag{5}$$

The OPEs between $N = 2$ currents and W-currents are

$$\begin{aligned}
J(z)W_2(w) &= \frac{2a^2(1+3a^2)}{(z-w)^3} + \frac{(1+3a^2)J(w)}{(z-w)^2} + \ldots, \\
J(z)W^+(w) &= \frac{(1+3a^2)G^+(w)}{(z-w)^2} + \frac{W^+(w)}{z-w} + \ldots, \\
J(z)W^-(w) &= \frac{(1+2a^2)G^-(w)}{(z-w)^2} + \frac{-W^-(w)}{z-w} + \ldots, \\
J(z)W_3(w) &= \frac{4a^2(1+3a^2)}{(z-w)^4} + \frac{(1+4a^2)J(w)}{(z-w)^3} + \frac{2W_2(w) + (1+2a^2)T(w)}{(z-w)^2} + \ldots,
\end{aligned} \tag{6}$$

$$\begin{aligned}
T(z)W_2(w) &= \frac{-2a^2(1+3a^2)}{(z-w)^4} + \frac{-(1+2a^2)J(w)}{(z-w)^3} + \frac{2W_2(w)}{(z-w)^2} + \frac{\partial W_2(w)}{z-w} + \ldots, \\
T(z)W^+(w) &= \frac{-(1+2a^2)G^+(w)}{(z-w)^3} + \frac{2W^+(w)}{(z-w)^2} + \frac{\partial W^+(w)}{z-w} + \ldots,
\end{aligned}$$

$$\begin{aligned}
T(z)W^{-}(w) &= \frac{-G^{-}(w)}{(z-w)^3} + \frac{3W^{-}(w)}{(z-w)^2} + \frac{\partial W^{-}(w)}{z-w} + \ldots, \\
T(z)W_3(w) &= \frac{-T(w)}{(z-w)^3} + \frac{3W_3(w)}{(z-w)^2} + \frac{\partial W_3(w)}{z-w} + \ldots,
\end{aligned} \tag{7}$$

$$\begin{aligned}
G^{+}(z)W_2(w) &= \frac{-W^{+}(w)}{z-w} + \ldots, \quad G^{+}(z)W^{+}(w) = \text{regular}, \\
G^{+}(z)W^{-}(w) &= \frac{4a^2(1+3a^2)}{(z-w)^4} + \frac{(1+4a^2)J(w)}{(z-w)^3} + \frac{2W_2(w)}{(z-w)^2} + \frac{W_3(w)}{z-w} + \ldots, \\
G^{+}(z)W_3(w) &= \frac{(1+4a^2)G^{+}(w)}{(z-w)^3} + \frac{2W^{+}(w)}{(z-w)^2} + \ldots, \\
G^{-}(z)W_2(w) &= \frac{a^2G^{-}(w)}{(z-w)^2} + \frac{W^{-}(w)}{z-w} + \ldots, \quad G^{-}(z)W^{-}(w) = \text{regular}, \\
G^{-}(z)W^{+}(w) &= \frac{2a^2(1+3a^2)}{(z-w)^4} + \frac{(1+2a^2)J(w)}{(z-w)^3} \\
&\quad + \frac{-2W_2(w) + a^2T(w)}{(z-w)^2} + \frac{W_3(w) - \partial W_2(w)}{z-w} + \ldots, \\
G^{-}(z)W_3(w) &= \frac{G^{-}(w)}{(z-w)^3} + \frac{3W^{-}(w)}{(z-w)^2} + \frac{\partial W^{-}(w)}{z-w} + \ldots .
\end{aligned} \tag{8}$$

Let us discuss the OPEs for W-currents. First we write down the OPEs between W_2 and $\{ W_2, W^{\pm}, W_3 \}$:

$$\begin{aligned}
W_2(z)W_2(w) &= \frac{1+4a^2-a^4-12a^6}{(z-w)^4} \\
&\quad + \frac{-2(1+a^2)W_2(w) + (1+2a^2)[(JJ) + 2a^2\partial J](w)}{(z-w)^2} \\
&\quad + \frac{-(1+a^2)\partial W_2(w) + (1+2a^2)[(\partial JJ) + a^2\partial^2 J](w)}{z-w} + \ldots,
\end{aligned} \tag{9}$$

$$\begin{aligned}
W_2(z)W^{+}(w) &= \frac{(1-4a^4)G^{+}(w)}{(z-w)^3} + \frac{-(1+a^2)W^{+}(w) + (1+2a^2)(JG^{+})(w)}{(z-w)^2} \\
&\quad + \frac{(JW^{+})(w) - (G^{+}W_2)(w) + (1+2a^2)(\partial JG^{+})(w) - \partial W^{+}(w)}{z-w} + \ldots,
\end{aligned} \tag{10}$$

$$\begin{aligned}
W_2(z)W^{-}(w) &= \frac{-(1+2a^2)G^{-}(w)}{(z-w)^3} \\
&\quad + \frac{-(1+a^2)W^{-}(w) + (1+2a^2)(JG^{-})(w) + a^2(1+2a^2)\partial G^{-}(w)}{(z-w)^2} \\
&\quad + \frac{\mathcal{O}^{2,-}_{(1)}(w)}{z-w} + \ldots,
\end{aligned} \tag{11}$$

$$\begin{aligned}
W_2(z)W_3(w) &= \frac{2(1+4a^2-a^4-12a^6)}{(z-w)^5}+\frac{(1+2a^2)J(w)}{(z-w)^4} \\
&+ \frac{-2(1+a^2)W_2(w)+(1+2a^2)[-a^2T+(JJ)+3a^2\partial J](w)}{(z-w)^3} \\
&+ \frac{\mathcal{O}^{2,3}_{(2)}(w)}{(z-w)^2}+\frac{\mathcal{O}^{2,3}_{(1)}(w)}{z-w}+\ldots,
\end{aligned} \tag{12}$$

where

$$\begin{aligned}
\mathcal{O}^{2,-}_{(1)}(w) &= -(JW^-)(w)+(G^-W_2)(w)-(1+a^2)\partial W^-(w) \\
&+(1+a^2)(\partial JG^-)(w)+a^2(J\partial G^-)(w)+a^2(\frac{1}{2}+a^2)\partial^2G^-(w), \\
\mathcal{O}^{2,3}_{(2)}(w) &= -W_3(w)+(JW_2)(w)-\partial W_2(w)-a^2(G^+G^-)(w)+(1+a^2)(JT)(w) \\
&+a^2\partial T(w)+(1+2a^2)[(\partial JJ)+\frac{3a^2}{2}\partial^2J](w), \\
\mathcal{O}^{2,3}_{(1)}(w) &= (G^+W^-)(w)+(G^-W^+)(w)+a^2(\partial G^-G^+)(w)+(\partial JW_2)(w)-\partial W_3(w) \\
&+(1+a^2)(\partial JT)(w)+(a^2+\frac{1}{2})(\partial^2JJ)(w)+a^2(a^2+\frac{2}{3})\partial^3J(w).
\end{aligned} \tag{13}$$

Here a normal ordered product $(AB)(z)$ for two generators $A(z)$ and $B(z)$ are defined as

$$(AB)(z)=\int\frac{dw}{2\pi\mathrm{i}}\frac{A(w)B(z)}{w-z}. \tag{14}$$

Second the OPEs among fermionic W-currents $W^{\pm}$ are given by

$$\begin{aligned}
W^+(z)W^+(w) &= \frac{-(1+2a^2)(G^+\partial G^+)(w)+2(G^+W^+)(w)}{z-w}+\ldots, \\
W^-(z)W^-(w) &= \frac{-(G^-\partial G^-)(w)-2(G^-W^-)(w)}{z-w}+\ldots, \\
W^+(z)W^-(w) &= \frac{2(1+4a^2-a^4-12a^6)}{(z-w)^5}+\frac{(1+2a^2)J(w)}{(z-w)^4} \\
&+\frac{(1+a^2-2a^4)T(w)-2(1+a^2)W_2(w)+(1+2a^2)[(JJ)+3a^2\partial J](w)}{(z-w)^3} \\
&+\frac{\mathcal{O}^{+,-}_{(2)}(w)}{(z-w)^2}+\frac{\mathcal{O}^{+,-}_{(1)}(w)}{z-w}+\ldots,
\end{aligned} \tag{15}$$

where

$$\mathcal{O}^{+,-}_{(2)}(w) = a^2W_3(w)-\partial W_2(w)+(JW_2)(w)+(1+a^2)(G^+G^-)(w)$$

$$\begin{aligned} & -a^2(JT)(w) - 2a^4\partial T(w) + (1+2a^2)[(\partial J J) + \frac{3a^2}{2}\partial^2 J](w), \\ \mathcal{O}^{+,-}_{(1)}(w) &= (JW_3)(w) - (TW_2)(w) + (\partial J W_2)(w) + a^2\partial W_3(w) - a^2(\partial T J)(w) \\ & +(1+a^2)(\partial G^+ G^-)(w) - a^4\partial^2 T(w) + (a^2+\frac{1}{2})(\partial^2 J J) + a^2(a^2+\frac{2}{3})\partial^3 J(w). \end{aligned} \tag{16}$$

Third the OPEs between $W^{\pm}$ and W_3 are

$$\begin{aligned} W^+(z)W_3(w) &= \frac{(1+2a^2)G^+(w)}{(z-w)^4} \\ & + \frac{-2(1+a^2)W^+(w) + (1+2a^2)[(JG^+) + (G^+J) + 3a^2\partial G^+](w)}{(z-w)^3} \\ & + \frac{\mathcal{O}^{+,3}_{(2)}(w)}{(z-w)^2} + \frac{\mathcal{O}^{+,3}_{(1)}(w)}{z-w} + \dots, \\ W^-(z)W_3(w) &= \frac{4(1+2a^2)G^-(w)}{(z-w)^4} + \frac{2(1+2a^2)\partial G^-(w)}{(z-w)^3} + \frac{\mathcal{O}^{-,3}_{(2)}(w)}{(z-w)^2} + \frac{\mathcal{O}^{-,3}_{(1)}(w)}{z-w} + \dots, \end{aligned} \tag{17}$$

where

$$\begin{aligned} \mathcal{O}^{+,3}_{(2)}(w) &= -\partial W^+(w) + (JW^+)(w) + (G^+W_2)(w) + (TG^+)(w) \\ & +(1+2a^2)[(\partial J G^+) + (\partial G^+ J)](w) + (-\frac{1}{2} + \frac{3a^2}{2} + 3a^4)\partial^2 G^+(w), \\ \mathcal{O}^{+,3}_{(1)}(w) &= [(G^+W_3) - (TW^+) + (\partial J W^+) + (\partial G^+ W_2)](w) + (1+a^2)(\partial G^+ T)(w) \\ & -a^2(\partial T G^+)(w) + (\frac{1}{2} + a^2)[(\partial^2 J G^+) + (\partial^2 G^+ J)](w) + (a^4 + \frac{2a^2}{3})\partial^3 G^+(w), \end{aligned} \tag{18}$$

and

$$\begin{aligned} \mathcal{O}^{-,3}_{(2)}(w) &= 2(JW^-)(w) + (G^-T)(w) + 2a^2\partial W^-(w) \\ & +(\partial G^- J)(w) - (\partial J G^-)(w) + 2a^2\partial^2 G^-(w), \\ \mathcal{O}^{-,3}_{(1)}(w) &= -(G^-W_3)(w) + (TW^-)(w) + (J\partial W^-)(w) + (\partial J W^-)(w) + (\partial G^- T)(w) \\ & +a^2\partial W^-(w) + \frac{1}{2}(\partial^2 G^- J)(w) - \frac{1}{2}(\partial^2 J G^-)(w) + \frac{2a^2}{3}\partial^3 G^-(w). \end{aligned} \tag{19}$$

Finally the OPE between W_3 and itself is found to be

$$W_3(z)W_3(w) = \frac{4(1+2a^2)T(w)}{(z-w)^4} + \frac{2(1+2a^2)\partial T(w)}{(z-w)^3} + \frac{\mathcal{O}^{3,3}_{(2)}(w)}{(z-w)^2} + \frac{\mathcal{O}^{3,3}_{(1)}(w)}{z-w} + \dots, \tag{20}$$

where

$$\begin{aligned}
\mathcal{O}^{3,3}_{(2)}(w) &= 2(JW_3)(w) - 2(G^+W^-)(w) + (TT)(w) - (\partial JT)(w) + (\partial TJ)(w) \\
&\quad +2a^2\partial W_3(w) + 2a^2\partial^2 T(w) + [(\partial G^+G^-) + (\partial G^-G^+)](w), \\
\mathcal{O}^{3,3}_{(1)}(w) &= \frac{1}{2}\partial\mathcal{O}^{3,3}_{(2)}(w) - \frac{1}{6}(1+2a^2)\partial^3 T(w). \qquad (21)
\end{aligned}$$

The equations (5)-(8) and (9)-(21) complete the operator product expansions of the $N=2$ super W_3 algebra. It is easy to see that the present definitions of the W-currents (4) give the Poisson algebra structure with quadratic relations in the classical limit of $a^2 \to -\infty$ [13].

Note that in the present definitions (3), the W-currents are quasi-primary fields with respect to the energy-momentum tensor $T^{N=2}(z)$. One may add suitable differential polynomials of the currents to these W-currents in order to get a $N=2$ supermultiplet $\{\widetilde{W}_2, \widetilde{W}^\pm, \widetilde{W}_3\}$ [3, 17]:

$$\begin{aligned}
\widetilde{W}_2 &= W_2 - \frac{a^2}{2}\partial J - \frac{3+8a^2}{2(5+18a^2)}(JJ) + \frac{(1-2a^2)(1+3a^2)}{5+18a^2}T^{N=2}, \\
\widetilde{W}^+ &= W^+ + \frac{(1-2a^2)(1+3a^2)}{5+18a^2}\partial G^+ - \frac{3+8a^2}{5+18a^2}(JG^+), \\
\widetilde{W}^- &= W^- - \frac{1+6a^2+12a^4}{5+18a^2}\partial G^- - \frac{3+8a^2}{5+18a^2}(JG^-), \\
\widetilde{W}_3 &= W_3 - \frac{1}{2}\partial W_2 + \frac{3+8a^2}{5+18a^2}[(G^+G^-) - (JT) + \frac{1}{4}\partial(JJ)] \\
&\quad -\frac{3+13a^2+18a^4}{2(5+18a^2)}\partial T^{N=2} - \frac{(2+3a^2)(1+2a^2)}{4(5+18a^2)}\partial^2 J. \qquad (22)
\end{aligned}$$

From (22) and the OPEs for the W-currents, we may compute the operator product expansions for the $N=2$ super-multiplet $\{\widetilde{W}_2, \widetilde{W}^\pm, \widetilde{W}_3\}$. After some calculations, one finds that these operator product expansions coincide with the results of the OPE method [14]. For example, one gets

$$\widetilde{W}_2(z)\widetilde{W}_2(w) = \frac{\frac{c^{N=2}\beta^2}{2}}{(z-w)^4} + \frac{\widetilde{\mathcal{O}}^{2,2}_{(2)}(w)}{(z-w)^2} + \frac{\frac{1}{2}\partial\widetilde{\mathcal{O}}^{2,2}_{(2)}(w)}{z-w} + \ldots, \qquad (23)$$

where $\beta = \sqrt{\frac{(1-2a^2)(2+3a^2)(1+4a^2)}{3(5+18a^2)}}$ and

$$\widetilde{\mathcal{O}}^{2,2}_{(2)}(w) = -\frac{6(1+2a^2)(1+5a^2)}{(5+18a^2)}\widetilde{W}_2(w) + \frac{2c^{N=2}\beta^2}{5+18a^2}[T^{N=2} - \frac{1}{4(1+3a^2)}(JJ)](w). \quad (24)$$

The relation between the present definitions and those in ref. [14] is given by formulae $(J, G^+, G^-, T^{N=2}) = (J, \sqrt{2}G, \sqrt{2}\bar{G}, T)$ and $(\widetilde{W}_2, \widetilde{W}^+, \widetilde{W}^-, \widetilde{W}_3) = -\beta(V, \sqrt{2}U, \sqrt{2}\bar{U}, W)$, where $(J, G, \bar{G}, T)$ and $(V, U, \bar{U}, W)$ are the $N=2$ primary W-currents in ref. [14].

It is known that the $A(n|n-1)$ Miura transformations give the Feigin-Fuchs representation of the $N=2$ CP_n super coset models[18] for $a^2 = -1/(k+n+1)$ [2, 3]. In these CP_n coset models there is a kind of duality relation such that $\{$ CP_n models at level $k=1$ $\} \equiv \{$ CP_1 models at level $k=n\}$ [18]. In the case of $n=2$, we get $a^2 = -1/4$. It easy to see that the W-currents $\widetilde{W}_2, \widetilde{W}^+, \widetilde{W}^-, \widetilde{W}_3$ have no central extensions and become zero-norm fields at this duality point[1]. Therefore we expect that these W-currents decouples from the representations of $N=2$ super-W_3 algebra[5]. But in order to check the decoupling we need to investigate the structure of the Verma modules more carefully, which is left for further investigation.

Although the present free field realization is obtained from the supersymmetric Miura transformation (2), the algebra does not depend on the particular free field realization of U_1, $G_1^{\pm}$ and T_1. This suggests the existence of other types of the free field realization of $N=2$ super-W_3 algebra, which would be useful for studying the quantum topological W_3-gravity theory.

The author would like to thank Wolfgang Lerche and Xiang Shen for useful discussions and the CERN Theory Division for its hospitality, where a part of this work was done.

---

[1] A similar phenomena has been observed in the case of Z_k parafermions[19]. The author would like to thank Wolfgang Lerche for pointing out this fact.

References

[1] T. Inami, Y. Matsuo and I. Yamanaka, Int. J. Mod. Phys. **A5** (1990) 4441;

[2] K. Ito, Phys. Lett. **B259** (1991) 73; Nucl. Phys. **B370** (1992) 123.

[3] D. Nemeschansky and S. Yankielowicz, USC preprint USC-91/005.

[4] K. Li, Phys. Lett. **B251** (1990) 54; Nucl. Phys. **B346** (1990) 329.

[5] M. Bershadsky, W. Lerche, D. Nemeschansky and N.P. Warner, preprint CERN-TH.6694/92.

[6] For review, see P. Bouwknegt and K. Schoutens, Phys. Rep. **223** (1993) 183.

[7] A.B. Zamolodchikov, Theor. Math. Phys. **63** (1985) 1205.

[8] V.A. Fateev and A.B. Zamolodchikov, Nucl. Phys. **B280** [FS18] (1987) 644; V.A. Fateev and S.L. Lukyanov, Int. J. Mod. Phys. **A3** (1988) 507.

[9] H.G. Kausch and G.M.T. Watts, Int. J. Mod. Phys. **A7** (1992) 4175.

[10] M. Bershadsky and H. Ooguri, Comm. Math. Phys. **126** (1989) 429.

[11] T. Inami and H. Kanno, J. Phys. **A25** (1992) 3729; K. Hiutu and D. Nemeschansky, Mod. Phys. Lett. **A6** (1991) 3179; J.M. Figueroa-O'Farill and E. Ramos, Nucl. Phys. **B368** (1992) 361; A. Das and W.-J. Huang, Mod. Phys. Lett. **A7** (1992) 2159.

[12] H. Lu, C.N. Pope, L.J. Romans, X. Shen and X.-J. Wang, Phys. Lett. **B264** (1991) 91.

[13] K. Ito, NBI preprint NBI-HE-92-77, to appear in the proceedings of the International Workshop "String Theory, Quantum Gravity and the Unification of Fundamental Interactions" Rome, September 21–26 (World Scientific, Singapore)

[14] L.J. Romans, Nucl. Phys. **B369** (1992) 403.

[15] S. Odake, Soryushiron Kenkyu (Kyoto) **78** (1989) 201 (in Japanese).

[16] J. Evans and T. Hollowood, Nucl. Phys. **B352** (1991)723; S. Komata, K. Mohri and H. Nohara, Nucl. Phys, **B359** (1991) 168.

[17] H. Lu, C.N. Pope, X.-J. Wang and K.W. Xu, Nucl. Phys. **B379** (1992) 24.

[18] Y. Kazama and H. Suzuki, Nucl. Phys. **B321** (1989) 232.

[19] F.J. Narganes-Quijano, Ann. Phys. **206** (1991) 494.